

C 100[®]

Anti-Stress for Fish, Shrimp & Poultry

Composition

Ascorbic Acid USP 100%

Indication

- To prevent vitamin C deficiency
- To manage various types of stress like: transportation, temperature change, water change, tank transfer, eye stalk ablation and any kind of disease stress
- To prevent stress after use of vaccines and antibiotics
- To increase the immunity of Fish, shrimp & poultry
- To increase the growth, survivability and hatchability of eggs
- To prevent hydrocarbon, heavy metal toxicity in water tank
- To de-chlorinate water

Dosage & Administration

For Fish & Shrimp

In water: 2-3 gm/ton

In feed: 1 gm/kg feed

For de-chlorination: 1gm **C 100** neutralizes (1mg of chlorine/liter of water) 380-liter water

For Poultry

In water: 1 gm **C 100** / 5-10 liter for 3-5 consecutive days or,

In feed: 1-3 gm **C 100** / 1000 kg feed for 3-5 consecutive days

Or, as directed by registered Veterinary/Aquaculture consultant

Direction of Use

Mix the required quantity of **C 100** with Nutrigel and mix with feed. Dry it for 10 min prior to use. In case of water Mix, the required quantity of **C 100** directly with water.

Contraindications

Not known.

Precautions

Once opened the pack should be used quickly. Store in a cool and dry place, away from light. Keep out of the reach of children.

Side effect

Not known.

Presentation

100g, 500g & 1 KG

For Aquaculture/Poultry Use Only